LINKS BETWEEN MATERNAL NUTRITION, MATERNAL COGNITION AND MENTAL HEALTH, AND CAREGIVING

Elizabeth Prado UC Davis Summit Institute of Development

Potential Pathways from Nutrition to Child Development



Adapted from Levitsky & Barnes (1972), Pollitt (1993), Prado & Dewey (2014)



Maternal Nutrition and Maternal Cognition

- Nutrition and brain function in adults
- Associations in women of reproductive age
 Iron-deficiency anemia (Murray-Kolb, 2011)
- □ 3 RCTs during pregnancy
 - Positive effect of iron supplements
 - 25 women in the US (Groner et al. 1986)
 - **81** women in South Africa (Beard et al. 2005)
 - Positive effect of multiple micronutrients (MMN) compared to iron/folic acid (IFA)
 - 640 women in Indonesia (Prado et al. 2012)



Maternal Nutrition and Maternal Depression/Anxiety

- Associations
 - Deficiencies in iron, zinc, B-vitamins and essential fatty acids have been associated with perinatal depression/anxiety (Leung & Kaplan 2009, Ramakrishnan 2011)
- 4 RCTs during pregnancy
 - Positive effect of iron supplements in IDA women in South Africa (Beard et al. 2005)
 - No effect of MMN vs IFA in Indonesia (Prado et al. 2012)
 - Positive effect of MMN vs IFA in Bangladesh (Frith et al. 2009)
 - Positive effect of MMN vs placebo in HIV-positive women in Tanzania (Smith Fawzi et al. 2007)



Supplementation with Multiple Micronutrients Intervention Trial (SUMMIT), Lombok, Indonesia

262 midwives randomly assigned to distribute a daily tablet to >30,000 women, containing:

<u>MMN</u> :	or	IFA: from	First visit to local pre-natal care	to 3 months post- partum				
lron Folic Acid	Niacin	Iron Folic Acid	clinic	parion				
Vitamin A	Vitamin B6 Vitamin B12							
Vitamin D Vitamin E Vitamin C Vitamin B1 Vitamin B2	Zinc Copper Selenium Iodine		I	birth				
		Cognitive Participants:						
		640 (344 MMN, 296 IFA) mothers tested after a mean of 25 WEEKS of supplementation (10% in the 1st or 2nd trimester, 39% in the 3rd trimester, 51% after birth).						

SUMMIT Study Group (2008). Effect of maternal multiple micronutrient supplementation on fetal loss and infant death in Indonesia: a double-blind cluster-randomised trial. *Lancet, 371*(9608), 215-227.

Test Selection Criteria (1) to (4) -

- 1) Tests that are designed to primarily tap aspects of specific cognitive and other brain-related functions, particularly those that have been associated with micronutrient deficiency in previous studies.
- 2) Tests that have been tied to particular brain structures and mechanisms, particularly those in which micronutrients play a role.
- 3) Tests that play a role in practical skills important for daily life.
- 4) Tests that are well-established and widely used.



Criteria (5) to (7) – Cross-cultural issues

- 5) Tests that are appropriate for the local setting.
- 6) Tests that are able to distinguish between different individual's abilities, so should result in a wide distribution of scores.
- 7) Tests with high reliability in the local setting.



Criteria (8) to (10) – Logistical issues

- 8) Tests that do not require extensive training for the testers (at maximum ~ 2-3 weeks), do not require previous experience in cognitive assessment, and do not require subjective judgments from the testers, such as the type of judgments a highly trained clinician would make.
- 9) Tests that are relatively brief to administer
- 10) Tests that do not require special equipment (only paper, pencil, stopwatch, and digital recorder).



Reliability and Validity

Test		Internal Reliability		Test-Retest Reliability		Correlation with education		
	n	Alpha/r	n	r	р	n	r	р
Set X								
Declarative Memory								
Word List Mem	ory							
Immediate R	ecall							
Delayed Rec	all 468	0.64	20	0.69	<.001	351	0.45	<.001
Delayed	468	0.68	20	0.76	<.001	347	0.42	<.001
Recognition	459	0.87	20	0.73	<.001	344	0.47	<.001
Working Memory								
Digit Span Forv	vard 467	0.67	20	0.89	<.001	350	0.31	<.001
Digit Span Bacl	kward 467	0.68	20	0.56	<.001	350	0.47	<.001
Set Y								
Motor Dexterity						·		
Coin Rotation								
Right Hand	0.07	0.04	20	0.69	<.001	244	0.23	<.001
Left Hand	307	0.61	20	0.44	.054	244	0.08	.211
Verbal Fluency	·							
Verbal Fluency	·	•	,					
Category: Fo	hod	0.44	11	0.96	<.001	243	0.09	.144
Category: Na	300	0.41	11	0.77	.005	243	0.16	.011
Depression Test (CES		0.83	21	0.84	<.001	240	-0.14	.028
Set Z	/		,					
Speed of Lexical Retri	eval							
Picture Naming			16	0.45	.082	414	0.42	<.001
Word Reading								
Real Words			15	0.95	<.001	353	0.53	<.001
Pseudoword	s		15	0.98	<.001	352	0.48	<.001
Mental Rotation Test	500	0.73	19	0.85	<.001	417	0.35	<.001

Effect of MMN on each Cognitive Motor, and Mood Test



Prado, E. L., et al. (2012). "The effect of maternal multiple micronutrient supplementation on cognition and mood during pregnancy and postpartum in Indonesia: a randomized trial." <u>PloS one **7**(3): e32519.</u>





□ Mean z-score across all cognitive tests



■IFA ■MMN

Maternal Cognition and Caregiving

- Positive associations between maternal cognition and
 - Infant and toddler dietary intake in Peru, Egypt, Kenya (Wachs et al. 2001, 2005; Bhargava et al. 2003)
 - HOME Inventory scores in Jamaica (Baker-Henningham et al. 2003)
 - Istening and reading comprehension of general health messages in Zambia (Stuebing et al. 1997)

Maternal Depression/Anxiety and Caregiving

- Association between maternal depression/anxiety and
 - Lower HOME Inventory scores in Jamaica (Baker-Henningham et al. 2003)
 - Reduced preference for breastfeeding in Barbados (Galler et al. 1999)
 - Use of health services in India (depressed mother were more likely to use health services than non-depressed mothers) Patel et al. (2002)

International Lipid-Based Nutrient Supplements (iLiNS) Project, Malawi

864 pregnant women randomized to 3 groups:







Adapted from Levitsky & Barnes (1972), Pollitt (1993), Prado & Dewey (2014)

Methods



- Maternal Depression/Anxiety
 - WHO Self-Reporting Questionnaire
 - Edinburgh Postnatal Depression
 Scale
 - Perceived Stress Scale
- Maternal Cognition
 - Digit Span Forward and Backward
 - Verbal Fluency
 - Mental Rotation
 - Functional Health Literacy

- Mother-Infant interaction
 - HOME Inventory
- Activity
 - Accelerometers
- Child Development
 - Motor: Kilifi Development Inventory (KDI)
 - Language: Adapted MacArthur-Bates Communicative
 Development Inventory
 - Socio-Emotional: Profile of Socio-Emotional Development (PSED)
 - Executive Function: A not B task

Thank you!

SUMMIT Team:

- Anuraj Shankar
- 🗖 Husni Muadz
- Michael Ullman
- Katie Alcock
- Mandri Apriatni
- Susy Sebayang
- Aditiawarman
- Ben Harefa
- Josephine Kadha
- Dini Prihatini
- Abas Jahari

iLiNS-Malawi Team:

- Kay Dewey
- Per Ashorn
- Ken Maleta
- Ulla Ashorn
- Steve Vosti
- John Phuka
- John Sadalaki
- Atupele Luwayo
- Rob Stewart
- Christine Stewart
- Anna Pulakka

SUMMIT Funding:

- U.S. State Department
 Fulbright grant
- National Science
 Foundation
- Turner Foundation
- UNICEF
- Centre for Health and Human Development
- USAID-Indonesia
- □ iLiNS Funding:
 - Bill & Melinda Gates
 Foundation